




MONOGRAPH · DOSING

Check menopause formulas against trial doses

You already compare forms and milligrams, and you want trial numbers next to label numbers. After this guide you will be able to parse any cycle or menopause panel, align each dose with its study, and price the daily serving with confidence.

- Dose first**
Match milligrams on the panel to milligrams in the trial.
- Form matters**
Elemental amount and extract detail decide if comparison works.
- Time trials**
Use twelve week logs and daily cost before judging response.

By Katherine Warren · Updated Sep 16, 2026 · 11 min read

CONTENTS

01	Start with the Supplement Facts panel	05	Calculate serving cost before committing
02	Map the usual ingredient categories	06	Take precise questions to the clinician
03	Identify blends that prevent comparison	07	The checklist
04	Monitor symptoms across eight to twelve weeks	08	Sources and further reading

By the final section you will be able to translate a cycle or menopause label into milligrams, compare those milligrams with study doses, and calculate what an eight to twelve week test truly costs. Two bottles can share a pink design and still deliver different quantities of each compound. One panel prints every amount, the other groups ingredients under one blend total, and only the first supports direct math. When the Supplement Facts panel becomes the reference, slogans carry less weight.

01

Start with the Supplement Facts panel

The capsule swallowed at night reflects agricultural, fermentation or marine sourcing decisions, yet the Supplement Facts panel is the sole location where those decisions turn into numbers. Federal rules direct that panel to open with the words Supplement Facts and to report serving size, servings per package, nutrient quantities and percent Daily Value. Dietary ingredients in the product must appear with names and quantities, together with serving size and servings per container.

That format counts because cross product comparison rests on amount per serving. A panel stating 100 mg magnesium per two capsule serving can be set beside dietary intake or a rival formula. The label puts magnesium inside a 900 mg blend while hiding the source amount. Labels list dietary ingredients in descending order of weight, which hints at what dominates a blend, though order alone never shows if any botanical reaches the level used in a trial.

Two minor points aid shelf review. Declaration of servings per container is not required when it equals the net quantity statement, for example 100 tablets where serving size is one tablet. Percent Daily Value indicates how a serving relates to daily intake levels, which assists when B vitamins or calcium with vitamin D join dietary sources. When a structure or function claim is used, such as supports immune health, labels must carry the statement that it has not been evaluated by the FDA and the product is not intended to diagnose, treat, cure or prevent any disease. That disclaimer neither confirms nor refutes worth. It identifies the claim as company language, not an agency conclusion.

Review the other ingredients line and any third party seal too, then revisit the central table. When names and quantities are whole, retain the bottle in the candidate group. When amounts are absent, remove it until the company supplies a certificate of analysis or a complete breakdown by lot and batch.

02

Map the usual ingredient categories

Cycle and menopause blends pull from a familiar pool, which permits label mapping component by component. Usual categories include magnesium, B vitamins, calcium with vitamin D, omega-3s, soy isoflavones, black cohosh and evening primrose oil. Dietary sources remain relevant here, because meals provide many of the same nutrients.

B vitamins serve as cofactors for enzymes that drive energy yielding metabolic pathways for carbohydrates, fats and proteins and support maintenance of nervous system functions. A menopause reference cites the recommended amount of vitamin B6 as 1.5 mg daily and vitamin B12 as 2.4 mcg daily. Inspect the panel for whether a formula meets, surpasses or misses those values, and recall that foods also contribute B6 and B12.

Magnesium recurs in midlife blends. Around half of people in the U.S. obtain too little magnesium from diet, and magnesium glycinate is a form that pairs magnesium with the amino acid glycine. The panel ought to declare the elemental amount plus the form. Lacking both, comparison against another product or against meals rich in beans, nuts and greens becomes difficult.

Calcium with vitamin D is often recommended for bone health during menopause, when sharp estrogen loss drives marked bone density loss and elevates osteoporosis risk. In a cross sectional study of 198 menopausal Korean women, dietary vitamin C intake was positively linked with femoral neck and total hip T-score measured by DEXA. During a 12 week interval before cognitive testing, 56 menopausal women in a prospective study took vitamin C at 50 mg daily or vitamin B12 at 0.5 mg daily. Those reports show associations and small trial designs, not evidence that a blend guards bone or cognition.

What sources say about modest or mixed effects

Soy isoflavones may lower hot flashes by about one fewer hot flash a day on average. Black cohosh shows mixed results across studies, with some reporting slight improvement in quality of life and hot flashes and recent studies reporting no difference from placebo. Soy, red clover and flaxseed phytoestrogens show mixed results for menopause symptoms and unclear effects on breast cancer risk. Evening primrose oil reviews describe efficacy for PMS and mastalgia in some trials but point to small samples, modest effects and need for regular use up to four to six months.

By contrast evening primrose oil holds linoleic acid at 60 percent to 80 percent and gamma linoleic acid at 8 percent to 14 percent. A 1994 pooled review of seven placebo controlled trials found a 3 to 6 g daily evening primrose oil dose tolerable in PMS volunteers. In a trial of 80 women with PMS, a daily dose of 1.5 g evening primrose oil for three months lowered symptom severity scores from 53.2 to 33.62, against 53.38 to 50.27 with placebo. Review authors state evening primrose oil should be used steadily for 4 or 6 months and an immediate response should not be expected. Those figures clarify why one month of use seldom resolves the question.

FROM THE EDITORS

Most women start it in their 40s, and perimenopause usually lasts around four years, though it can last for only a short time or stretch for up to a decade. A bottle purchased for a brief transition is judged on a different timeline than one taken across several years, which makes dated notes valuable.

03

Identify blends that prevent comparison

A proprietary blend reports a combined weight for multiple ingredients while withholding each dose. The descending order by weight rule remains in force, so the first constituent exceeds the second, although the margin could be wide or narrow. Whether a specific multi ingredient blend supplies an effective dose of each listed herb or nutrient when amounts are hidden in proprietary blends is not shown in the evidence reviewed here.

That lack of detail defeats the two comparisons label readers seek. One is dose compared with study intake, for instance 1.5 g evening primrose oil or 50 mg vitamin C used in studies. The other is formula versus formula. Two labels may each show black cohosh, soy and magnesium while holding quite different quantities. Absent individual amounts, extract ratio or standardisation details, and lot level paperwork, the purchaser cannot determine which option reflects the supply chain narrative on the front label.

Melissa Tharpe, womens health nurse practitioner, states the cost in plain terms: Heavily advertised supplements with a blend of ingredients are probably a waste of money. Her method advice is equally plain: "I always suggest trying one supplement at a time so you can see how your body responds." Testing one product at a time aids isolation of response. Which ingredient in a blend accounts for any symptom shift without testing one ingredient at a time remains unknown, and mixed blends keep that question unresolved.

Request the materials that permit comparison from manufacturers. A certificate of analysis from a third party lab can verify identity and screen for heavy metals. A lot number ties the bottle to its harvest and its encapsulation run. Standardisation statements and fill weight define

what the extract was intended to supply. When a brand can identify growers and present paperwork by batch, acknowledge that openness. When that stage stays opaque, log it as opaque.

04

Monitor symptoms across eight to twelve weeks

Cycle and menopause symptoms vary with week, stress and season, so recollection alone frequently misguides. Written tracking across eight to twelve weeks yields a more stable record than memory at a follow up visit. How long anyone needs to try a formula before judging response beyond the three month and twelve week periods used in trials is not established, but those trial windows give a practical point to review.

Select a tool that will truly see daily use. Options include an ACOG tracker from a professional body, a phone app or a paper notebook kept beside the bottles. Daily entries perform best, even when short. Productive items to log include bleeding changes, hot flash count and minutes, sleep hours and interruptions, mood, vaginal and urinary changes, memory lapses and pain. Record serving size taken, time of day and any missed days. Record illness, travel, alcohol, new medicines and menstrual day where relevant.

Daily symptom tracking lowered reported symptoms and loneliness in a two week study, but authors note simply being in a study and knowing others share experiences may account for part of the effect. One study participant described that shared effect in simple terms: "I feel more normal that other women are doing the same survey and are probably experiencing similar issues, especially the emotional and mental ones." Anticipate some early shift from attention by itself. Extended records, maintained the same way each day, simplify separation of normal variation from a pattern worth raising.

FROM THE EDITORS

Keep a current list of medicines and supplements with doses, and bring it to every visit. Whether lasting use of herbal blends is safe for people with hormone sensitive conditions or on blood thinners and other medicines is not settled.

05

Calculate serving cost before committing

Price per bottle deceives when serving sizes diverge. Cost per serving says much about the field to capsule chain, from grower pay and extraction to testing, encapsulation and retail margin. For calculation, divide bottle price by servings per container. Then adjust for the amount in each serving. A bottle that seems expensive may prove moderate by the day, while an inexpensive bottle dosed at four capsules daily may prove dearer.

Apply the panel to the arithmetic. Locate serving size, for example two capsules, and servings per container. Divide to obtain daily cost at suggested use. If the plan covers a twelve week review, multiply daily cost by 84 days. That sum frames the self test the way researchers frame theirs. Twelve week trials for vitamin C and three month trials for evening primrose oil define a review window, and review authors for evening primrose oil cite four to six months before judging.

The argument for pausing before a large cocktail is practical as well as financial. Multi ingredient products bundle several unknowns at a single price. When individual doses stay hidden, the buyer funds constituents that may occur in tiny amounts. That worry fits our habit of checking the label against the trial before paying for hyped blends. A simpler label with complete doses is often cheaper to test and produces cleaner notes.

Retain receipts and lot numbers alongside the symptom log. If a product alters colour, smell or fill weight between bottles, note the lot and ask the maker whether grower, harvest or extraction changed. Place names, dates and quantities outweigh front label terms, and words like pure or natural require a document behind them to carry meaning.

06

Take precise questions to the clinician

Thorough notes improve a visit. Carry the symptom record, the bottles or legible photos of Supplement Facts panels, the daily cost and the dates used. Name the targeted symptom or issue, note what the maker states and what the label shows. Ask what the evidence base suggests, including that supplement benefits look small and mixed beside prescription options.

Ask about safety with personal conditions and medicines, including hormone sensitive conditions, blood thinners and other prescriptions. So ask about other paths, including lifestyle steps and prescription options. Systemic estrogen in pill, patch, spray, gel or cream form is called the best treatment for hot flashes and night sweats, while vaginal estrogen releases a small amount for dryness, pain during sex and some bladder symptoms. Fezolinetant and elinzanetant are hormone free medicines that treat menopause hot flashes by blocking one or

more brain pathways that help control body temperature. Certain antidepressants called selective serotonin reuptake inhibitors may soften hot flashes in menopause. Those choices hold their own benefits and risks, which a clinician can balance with history and preferences.

Report certain shifts promptly rather than holding them for the twelve week review. Ask the clinic which bleeding changes, mood shifts, urinary symptoms or pain should prompt a call. Perimenopause can extend across years, and plans may call for revision as bleeding patterns, sleep and priorities shift. A concise written question set holds the visit on track: what is the target, what has shifted in the log, is the current product transparent enough to judge, should use continue, pause or switch, and what would indicate need for a different approach.

07

The checklist

-
- ☐ Snap the Supplement Facts panel plus other ingredients text
 - ☐ Copy serving size and servings per container figures
 - ☐ Transcribe every dose with percent Daily Value
 - ☐ Flag every proprietary blend that keeps amounts hidden.
 - ☐ Assess one supplement at a time where feasible
 - ☐ Chart bleeding, hot flashes, sleep, mood plus pain each day
 - ☐ Log intake time, skipped days and cycle day
 - ☐ Work out daily price and 84 day total
 - ☐ File lot numbers and receipts beside the log
 - ☐ Carry full medicine and condition inventory to visits
 - ☐ Query proof, safety and prescription substitutes
 - ☐ Fix a reassessment date at eight to twelve weeks
-

Sources and further reading

- Selected vitamins and quality of life in menopausal women · [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)
- Evening Primrose (Oenothera biennis) Oil in Management of Female Ailments · [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)
- Dietary Supplement Labeling Guide: Chapter IV. Nutrition Labeling · [fda.gov](https://www.fda.gov/)
- Perimenopause - Diagnosis and treatment · [mayoclinic.org](https://www.mayoclinic.org/)
- Should you take supplements during perimenopause and menopause? · [novanthealth.org](https://www.novanthealth.org/)
- How tracking menopause symptoms can give women more control over their health · theconversation.com
- Menopause Symptom Tracker · [acog.org](https://www.acog.org/)

For information only, not medical advice. These statements have not been evaluated by the Food and Drug Administration, and no supplement here is intended to diagnose, treat, cure or prevent any disease. Ask a doctor or pharmacist before starting one, especially alongside prescription medicines, in pregnancy or before surgery.



WRITTEN BY

Katherine Warren

Katherine edits the news desk and the monographs at Half Life. She keeps a spreadsheet of every dose she has seen on a label beside the dose used in the trial, and the gap between the two columns is usually the story.